



The GUD technique: glandar urethral disassembly. An alternative for distal hypospadias repair

Antonio Macedo Jr. ^{1,2}, Sérgio Leite Ottoni ¹, Gilmar Garrone ¹, Riberto Liguori ¹, Ricardo Marcondes Mattos ¹, Marcela Leal da Cruz ¹

¹ Departamento de Urologia, CACA-NUPEP, São Paulo, SP, Brasil; ² Departamento de Pediatria, Universidade Federal de São Paulo, São Paulo, SP, Brasil

ABSTRACT

Introduction: We present an alternative procedure for distal hypospadias consisting of urethral mobilization and partial glandar disassembly, namely GUD (glandar urethral disassembly) technique.

Materials and Methods: A subcoronal circumcision exposes distal dysplastic urethra. We incise the Buck's fascia on both sides of urethra releasing it partially from the corpora. We keep a thin bridge of urethral plate to the glans and disassembly almost completely the glans from the corpora, except for the bridge. The glans is incised creating two wide wings that are extremely mobile. The urethra is mobilized, advanced and sutured to the tip of the glans. The glans wings embrace the distal urethra producing a conical glans.

Discussion: The concept of urethral mobilization has been reported and popularized by Koff in the literature to correct distal hypospadias. One of the limitations of this procedure is the risk of urethral retraction due to extensive proximal dissection. We got inspiration from Mitchell and Bagli's work of penile disassembly in epispadias to develop the GUD concept. We adopt minimal urethral mobilization mainly in glandar/proximal penile shaft and complete deconstruction of the glans, detaching the corpora from the glans and rotating the wide glans wings to embrace the urethra. Therefore we avoid suture urethroplasty and refurbish the glans to a better conical shape.

Conclusion: We are convinced that this operation can be regarded as a genuine alternative to distal hypospadias (coronal and subcoronal) but should not be addressed to midshaft forms.

ARTICLE INFO

 **Marcela Leal da Cruz**

<http://orcid.org/0000-0002-7087-4534>

Keywords:

Hypospadias; Urethra;
Circumcision

Int Braz J Urol. 2020; 46: 1072-4

Submitted for publication:
March 13, 2019

Accepted after revision:
June 09, 2019

Published as Ahead of Print:
August 20, 2019

INTRODUCTION

Distal hypospadias can be treated by different techniques although TIP repair has been regarded as the preferred choice by many authors (1). Systematic literature review reports urethroplasty complications in approximately


7% of distal TIP repairs, even though other authors report higher complications rate (2).

In this video we want to describe an alternative procedure for distal hypospadias consisting of urethral mobilization and partial glandar disassembly (the GUD technique).

MATERIALS AND METHODS


A subcoronal circumcision is performed and the entire penis shaft is degloved exposing distal urethra. We incise the Buck's fascia on both sides lateral to distal urethra releasing it from the corpora. From this standpoint, we dissect the glans and disassembly it almost completely from the tip of corpora, except for a bridge of tissue similar to urethral plate connecting the urethra to the tip of glans. The urethra is mobilized and the glans incised creating two very mobile wings that allow a thorough refurbishment of the glans (Figure-1). No major bleeding from the spongy tissue occurs when the disassembly process respects the anatomy and the penis is compressed with a tourniquet on its base.

Figure 1 - GUD Technique – Partial Glandar Disassembly.



We suture the urethra to the glans with interrupted 7.0 PDS sutures. The spongy tissue of the distal urethra is sutured in the midline with 6.0 PDS. The lateral borders of the glans are brought together over the urethra with subcutaneous/subepithelial 5.0 PDS sutures concluding the glans reconstruction. A 6F silicone catheter is left indwelling for 7 days just to avoid voiding discomfort in case of glans edema (Figure-2). A

Figure 2 - GUD technique: final aspect.



Tegaderm dressing is applied to the shaft of the penis concluding the procedure.


DISCUSSION

Koff et al. published a modification of the Barcat technique known as extensive urethral mobilization and confirmed excellent cosmetic and functional results on 168 patients with only 3.5% of the patients requiring reoperation (3). This technique has been also reproduced by several other groups.

Mitchell & Blagi (4) and Perovic et al. (5) reported on complete penile disassembly for epispadia repair as a way to complete release of the rotation of the penis and treat dorsal chordee bringing the urethra to a more functional location. We combined both procedures to propose the GUD technique. The rationale for this procedure is to avoid suture urethroplasty, except sutures to reposition the urethra distally and without tension. The aggressive partial glandular disassembly is the major difference compared to other modifications of Koff's technique, that rely mostly in proximal

mobilization of the urethra up to the bulbar area. Those procedures incise the glans to get two wings but do not disconnect it from corpora totally. We are convinced the GUD technique combines up-mobilization of the urethra with down mobilization and rotation of glans wings and the result is a conical and better cosmetic glans (Figure-3).

Figure 3 - GUD technique – Detail of conic aspect of the glans.



CONCLUSIONS

We are convinced that this operation can be regarded as a genuine alternative to distal hy-

pospadias (coronal and subcoronal), but do not recommend it for midshaft forms.

CONFLICT OF INTEREST

None declared.

REFERENCES

1. Snodgrass W. Tubularized, incised plate urethroplasty for distal hypospadias. *J Urol.* 1994;151:464-5.
2. Wilkinson DJ, Farrelly P, Kenny SE. Outcomes in distal hypospadias: a systematic review of the Mathieu and tubularized incised plate repairs. *J Pediatr Urol.* 2012;8:307-12.
3. Koff SA, Brinkman J, Ulrich J, Deighton D. Extensive mobilization of the urethral plate and urethra for repair of hypospadias: the modified Barcat technique. *J Urol.* 1994;151:466-9.
4. Mitchell ME, Bāgli DJ. Complete penile disassembly for epispadias repair: the Mitchell technique. *J Urol.* 1996;155:300-4.
5. Perovic SV, Vukadinovic V, Djordjevic ML, Djakovic NG. Penile disassembly technique for epispadias repair: variants of technique. *J Urol.* 1999;162(3 Pt 2):1181-4.

Correspondence address:

Marcela Leal da Cruz, MD
 Departamento de Urologia,
 CACAU-NUPEP
 Rua Maestro Cardim, 560/215
 São Paulo, SP, 01323-000, Brasil
 Fax: +55 11 3287-3954
 E-mail: ma_celaleal@yahoo.com.br